



From the journal:  
**Chemical Communications**  
 14 November 2017, Issue 88.

# Improving Na–O<sub>2</sub> batteries with redox mediators

James T. Frith,<sup>a</sup> Imanol Landa-Medrano,<sup>bc</sup> Idoia Ruiz de Larramendi,<sup>b</sup> Teófilo Rojo,<sup>bd</sup> John R. Owen<sup>a</sup> and Nuria Garcia-Araez<sup>\*a</sup>

⊖ Author affiliations

\* Corresponding authors

<sup>a</sup> Department of Chemistry, University of Southampton, Southampton, UK  
 E-mail: [n.garcia-araez@soton.ac.uk](mailto:n.garcia-araez@soton.ac.uk)

<sup>b</sup> Departamento de Química Inorgánica, Facultad de Ciencia y Tecnología, Universidad del País Vasco (UPV/EHU) Barrio Sarriena s/n, 48940 Leioa, Spain

<sup>c</sup> BCMaterials, Parque Científico y Tecnológico de Bizkaia, 48160 Derio, Spain

<sup>d</sup> CIC EnergiGUNE, Albert Einstein 48, Miñano, Spain

## Abstract

A new route to enhance the performance of Na–O<sub>2</sub> cells is demonstrated. Redox mediators (such as ethyl viologen) are shown to facilitate the discharge reaction, producing an increased capacity (due to suppressed electrode passivation), higher discharge potential (due to faster kinetics) and stable cycling.

